

# Assessing the impacts of ocean acidification upon tropical tuna

*Don Bromhead, Vernon Scholey, Daniel Margulies, Simon Nicol, John Wexler, Maria Santiago, Simon Hoyle, Cleridy Lennert-Cody, John Havenhand, Jane E. Williamson, Tatiana Ilyina and Patrick Lehodey\**

## Introduction

Increasing concentrations of CO<sub>2</sub> in the Earth's atmosphere (IPCC 2007) are causing a gradual warming and acidification of the Earth's oceans (e.g. Barnett et al. 2005; Caldeira and Wickett 2003; Feely et al. 2004). Both warming and acidification have the potential to affect the distribution and population dynamics of many marine organisms (IPCC 2007; Raven et al. 2005; Fabry et al. 2008). Significant advances in knowledge have been made over the last decade that have advanced understanding of how increasing ocean acidity will impact nearshore and coral reef ecosystems (Fabry et al. 2008). Our understanding about the effects of acidification on pelagic ecosystems, however, remains rudimentary. In the Pacific Ocean, improving our knowledge on the possible impacts on the pelagic environment is important, as the Pacific's tuna populations are of one of the largest and most valuable fisheries in the world (Williams and Terawasi 2009). The income derived from tuna fisheries provides a significant contribution to the economies of many Pacific Island countries and territories (Gillett 2009). To ensure such economic benefits are maintained through the sustainable management of this fishery requires an understanding of not only fishery impacts, but impacts of other factors upon population biomass and structure over time. While fishery scientists are now attempting to predict how ocean warming will affect Pacific tuna populations (Lehodey et al. 2010, 2013), no one has previously investigated how ocean acidification (OA) may affect these species and associated fisheries.

To advance our knowledge of the impacts of OA upon tuna populations and fisheries, a pilot study was undertaken at the Inter-American Tropical Tuna Commission (IATTC) Achotines Laboratory in Panama. The objectives of the pilot study were to develop and test experimental protocols to examine the potential effects of OA on yellowfin tuna (*Thunnus albacares*) egg fertilisation, egg and larval development, growth, and survival, and on the rapid selection of resistant genotypes. The following article provides an overview of the project, including a description of the trials conducted and a summary of the results. A full description of the study and the results has been submitted to *Deep Sea Research Part II* for publication.

## Projected changes in ocean acidity

Concentrations of CO<sub>2</sub> in the ocean (pCO<sub>2</sub>) tend towards equilibrium with the CO<sub>2</sub> in the atmosphere. Since the start of the industrial revolution, the world's oceans are estimated to have absorbed about 30%–50% of global man-made CO<sub>2</sub> emissions, which has lowered the average sea-surface pH by 0.1 units (i.e. making the ocean more acidic and less alkaline) (Feely et al. 2004; Sabine et al. 2004; Orr et al. 2005). It is estimated that uptake of atmospheric CO<sub>2</sub> by global oceans will further reduce sea-surface pH by 0.3–0.4 units by 2100 and up to 0.7 units by the year 2300 (Caldeira and Wickett 2003, 2005).

Concentrations of pH show spatial heterogeneity both between oceanic regions (surface waters) and throughout the water column. Currently, surface layer pH values are lowest in higher latitudes and areas where upwellings may bring subsurface waters with lower pH to the surface. Although the mean seawater pH is expected to decrease globally, the rate of this change is predicted to be greater in high latitudes, and lower in tropical and subtropical waters (Ilyina et al. 2013; Bopp et al. 2013). The degree of change is also dependent on future anthropogenic CO<sub>2</sub> emissions and whether these are higher or lower than “business as normal” IPCC projections, and may be impacted by other predicted consequences of climate change, such as changed ocean circulation patterns (IPCC 2011).

In the Pacific Ocean there is considerable seasonal and vertical/horizontal spatial variation in pH and pCO<sub>2</sub>. In the eastern tropical Pacific Ocean, surface water pH is on average lower than in the western tropical Pacific, and the pH at 50 m depth (in the eastern Pacific) is on average 0.54 pH units less than at the surface. Across the region of tuna spawning habitat, mean surface water pH is predicted to decrease between 0.26 and 0.49 pH units by 2100 (under the high atmospheric CO<sub>2</sub> scenario of IPCC) (Ilyina et al. 2013). In the eastern Pacific Ocean the pH of surface waters is predicted to decrease from 8.05 to about 7.73 while in the western Pacific Ocean, the mean decline in pH is projected to be 0.40 pH units (with a maximum predicted decline of 0.46) (Ilyina et al. 2013).

\* Author affiliations are given at the end of the article.

---

## Impacts on fish

Our synopsis of the available literature suggests that the early life stages of fish may be more vulnerable to direct OA impacts than adult fish, due to different modes of respiration and ion exchange (Jonz and Nurse 2006; Pelster 2008). Adult fish generally show a stronger capacity to compensate for prolonged exposure to elevated pCO<sub>2</sub> due to their ability to control acid base balance by bicarbonate buffering across the gills and to a more limited degree, via the kidneys (Brauner and Baker 2009; Esbaugh et al. 2012). Conversely, exposure to elevated pCO<sub>2</sub> has been found to adversely affect embryonic development (Tseng et al. 2013), larval and juvenile growth (Baumann et al. 2011), tissue/organ health (Frommel et al. 2011), and survival (Baumann et al. 2011). Such effects however appear to be species specific, with numerous studies failing to detect direct relationships between “near future” levels of elevated pCO<sub>2</sub> and embryogenesis (Franke and Clemmessen 2011), hatching (Frommel et al. 2012), growth and development (Bignami et al. 2013a; Hurst et al. 2012, 2013; Munday et al. 2011a; Frommel et al. 2011, 2012), swimming ability (Bignami et al. 2013a; Munday et al. 2009a) or survival (Frommel et al. 2012; Munday et al. 2011a). Similar results are reported for the growth of otoliths, with increased size and/or density observed in some species (Checkley et al. 2009; Munday et al. 2011b; Hurst et al. 2012; Bignami et al. 2013a, 2013b) but not in others (Frommel et al. 2012; Franke and Clemmessen 2011; Munday et al. 2011a).

The sub-lethal effects of OA may present the largest risk to individuals and populations for many species (Briffa et al. 2012). Somatic impacts such as reduced growth and size at age have been linked to increased mortality in natural fish populations due to increased risks of predation and reduced ability to find food (e.g. Houde 1989; Leggett and Deblois 1994). Otoliths play an important role in detecting sound, acceleration and body position in fish (Bignami et al. 2013b) and altered otolith formation may change predator avoidance and foraging efficiencies of individuals. Neurological and behavioral effects associated with elevated CO<sub>2</sub> levels have also been observed in coral reef fish (Briffa et al. 2012). These include disruption to mating propensity (Sundin et al. 2013), impaired ability to make settlement choices (Munday et al. 2009b), altered timing of settlement (Devine et al. 2012), impaired response to predator and prey olfactory cues (Allen et al. 2013; Cripps et al. 2011; Dixon et al. 2010; Nilsson et al. 2012), reduced escape distances (Allen et al. 2013), altered responses to visual threats (Ferrari et al. 2012a) and auditory signals (Simpson et al. 2011), behavioral lateralisation (Domenici et al. 2012; Nilsson et al. 2012) and a reduced capacity to learn (Ferrari et al. 2012b). Such behavioral changes were recently linked in reef species to the impact of

elevated pCO<sub>2</sub> upon key brain neurotransmitter GABA-A (Nilsson et al. 2012), with potential implications for other species given the highly conserved nature of GABA-A across species.

## Potential vulnerability of tunas

While many of the original studies in this area focused on reef species, recent studies have expanded to include non-reef species (e.g. Bignami et al. 2013a, 2013b; Frommel et al. 2010, 2011, 2012; Sundin et al. 2013; Tseng et al. 2013), but these have not included tuna species. The only experiment known to have tested impacts upon a tuna species (eastern little tuna) tested pCO<sub>2</sub> levels that far exceeded those predicted using IPCC scenarios (Kikkawa et al. 2003). Pelagic fish species are considered in general to have evolved in a relatively more stable pH environment than coastal and reef species, and this, in addition to low blood pCO<sub>2</sub> levels that can be expected to result from their high rates of gas exchange, may make them more vulnerable to changes in ocean pCO<sub>2</sub> and pH (e.g. Nilsson et al. 2012).

Tropical tunas range and spawn in equatorial and subtropical waters from the far western Pacific Ocean to the far eastern Pacific Ocean, and in both near coastal waters and oceanic waters (Schaefer 2001). It is unknown whether the historical level of OA variability across the tropical Pacific has been sufficient to have conferred some evolved resilience in tropical tunas to predicted future levels of OA.

Although the likely effects of OA on tuna populations have not been investigated, such research is clearly a high priority. Decision-makers need timely and appropriate scientific advice for current and future tuna fisheries management and worldwide adaptation planning. In October 2010 the Pelagic Fisheries Research Program (PFRP) recognised this need and funded a collaborative study of the impact of OA on the early developmental stages of Pacific yellowfin tuna. The study, led by the Secretariat of the Pacific Community and IATTC, is investigating the effect of OA upon embryonic development, hatching rates, condition, development, and growth and survival in pre- and post-feeding yellowfin larvae. The collaboration includes scientists from the Max Planck Institute of Meteorology (Germany), Collecte Localisation Satellites (France), the University of Gothenburg (Sweden), and Macquarie University (Australia). As tolerance to OA has been found to be variable on an individual level in other species, the project has included a component to look at whether genotypes (the genetic makeup) of individual yellowfin larvae vary in their responses to different CO<sub>2</sub> levels. This last component is a first step towards determining if OA causes genetic selection of

resistant genotypes in this species. Analyses to estimate the effects of OA on life stage development and rates of deformity, otolith formation and genetics are ongoing and not presented.

## Achotines facility

Experimental trials were conducted at IATTC's Achotines Laboratory, Panama, in October and November 2011. The facility was inaugurated in 1985 and is one of only a few in the world with the location, equipment, and expertise to conduct investigations of this type. Achotines Bay is located on a section of coastline (Fig. 1) where the continental shelf drops rapidly and deep oceanic waters are close to shore, allowing researchers to easily access local tuna populations for either field studies of early development or to obtain yellowfin tuna as captive broodstock. At the lab an in-ground concrete tank holds a broodstock population of yellowfin tuna that have spawned on a near-daily basis since October of 1996, providing a reliable source of eggs and larvae for early developmental studies.

## Experimental trials

Experimental trials were grouped into three categories: sperm and fertilisation trials, egg and larval trials, and genetics analyses. The egg and larval trials were

replicated, with the initial trials in October and the replications in November.

## Tank set up and pH control

The experimental setup for the fertilised egg and larval trials consisted of 15 experimental tanks. Each 840-liter (l) capacity tank was nested inside of an 1100-l tank filled with seawater that acted as a buffer/insulator to stabilise the water temperature in the smaller tank during the trials. Water flow, lighting, aeration, and turbulence levels were adjusted to set these parameters as uniformly as possible across all tanks.

Four pH treatment levels (6.9, 7.3, 7.7 and 8.1) were targeted in each trial with three replicate tanks per treatment level (Fig. 2). These levels were chosen based on results from the ocean-carbon-cycle models using the IPCC IS92a Scenario (e.g. the Hamburg Ocean Carbon Cycle model) (Ilyina et al. 2009) and reference to published studies on predicted pH levels (Ilyina et al. 2013; Caldeira and Wickett 2003) and take into account spatial variation in predicted declines, not just the global average predicted declines. Target pH levels 6.9 to 8.1 encompass potential mean ocean pH levels estimated for the current oceans and predicted for near future oceans (to 2300). A fifth treatment with a pH of 6.5 was also included. This treatment was applied as a range test to ascertain that extreme values of pH would result in mortality. This value is well below the lowest predicted pH for the time period to 2300.




Figure 1. The location of the Achotines Laboratory. The bay in the centre of the photo is Achotines Bay.



Figure 2. The multiple experimental tanks with nested egg incubators used in experiment 1 and experiment 2 of the pilot study.

The local coastal waters that supply the Achatines Laboratory seawater system were very close to pH 8.2 during the trial period and therefore ambient seawater was used for the high pH level. The four lower treatment levels of seawater pH were maintained by regulation of mixtures of compressed air and CO<sub>2</sub> bubbled through air diffusers in each tank. The use of CO<sub>2</sub> was critical to modifying water chemistry (i.e. increasing carbonic acid, increasing H<sup>+</sup>, lowering pH) in a manner consistent with CO<sub>2</sub>-induced ocean acidification.

In all trials, water-quality parameters (pH, temperature, salinity, dissolved oxygen, CO<sub>2</sub>, alkalinity) were measured at frequent intervals in each tank. Controlling pH in ocean-acidification experimental systems is a difficult task (Riebesell et al. 2010). For these experiments, sophisticated electronic gas-flow controllers were used to precisely control the mix of air and CO<sub>2</sub> supplied to the tanks in each module. The average pH attained in each module (treatment level) for the first experiment was within 0.15 units of the target pH (and generally much closer). In the second experiment average pH levels in each module varied, at times, by several units from the target pH.

### Fertilised egg and larval trials

The effects of OA upon mortality, growth, and development of eggs and larvae of yellowfin tuna were tested by rearing larvae from egg stage to first-feeding stage in

15 tanks comprising the five treatment (pH) levels. Trials were continuous but effectively comprised three phases: egg phase, yolk-sac larval phase, and first-feeding larval phase, with sampling regimes differing in each phase.

To start the experiment, fertilised eggs were collected from a daily spawn in the broodstock tank of the Achatines Laboratory and randomly stocked in each of 15 cylindrical egg-incubation nets nested one per experimental tank (Fig. 3). Eggs were stocked in each egg-incubation net at a density of 177 eggs L<sup>-1</sup>, and eggs were immediately sampled fresh for weights and measurements. Additional samples of eggs were taken during the incubation period and fixed for subsequent histological examination of tissue and organ development.

Yolk-sac larvae were then dispersed from the egg-incubation nets into their respective experimental tanks 1 hour (h) after hatching. The yolk-sac phase in yellowfin tuna larvae continues until approximately 50–70 h after hatching depending on water temperature. Yolk-sac-stage and feeding larvae were maintained in the same experimental tanks until early on the seventh day of feeding (approximately 8.5 days post-hatching; 9.38 days total from egg transfer), when the experiments were terminated.

Larvae were fed cultured *Brachionus plicatilis* (rotifers) at densities of 3–5 mL<sup>-1</sup>. Dense blooms of unicellular algae (500,000–750,000 cells mL<sup>-1</sup>) were maintained in each tank to facilitate rearing (Margulies et al. 2007).




Figure 3. The cylindrical egg-incubation nets used in the pilot study. One net was nested in each experimental tank.

During each phase (egg, yolk-sac, and first-feeding), 15–17 fresh samples were taken at various intervals from each tank to be measured (total length, notochord length, body depth at pectoral, body depth at vent) and processed for dry-weight determination. Samples collected during each developmental phase were also fixed and stored for later analyses of tissue histology and organ development, feeding success, genetic variability and otolith development.

At the termination of the experiment each tank was slowly drained of water and all surviving larvae were removed by beaker and counted. The percentage of expected survival (adjusted for sample removals) was estimated for each tank.

## Results and discussion

Our study tested the effect of a range of pH (~6.9–8.2) and pCO<sub>2</sub> (330–10,467 microatmospheres [uatm]) conditions upon yellowfin tuna eggs and larvae. This range is broad enough to take into account current and future (to the year 2300) spatial variability in water chemistry across the yellowfin tuna spawning habitat range in the Pacific. Experimental tanks with mean pCO<sub>2</sub> of less than 2500 uatm provided conditions that are relevant to the assessment of “near future” (i.e. year 2100) impacts on yellowfin tuna. Over longer time frames (e.g. to 2300) or with further increases in CO<sub>2</sub> emissions, pCO<sub>2</sub> may increase further and pH decrease further (e.g. by

0.7 pH units according to Caldeira and Wickett [2003]), in which case our experimental tanks with mean pCO<sub>2</sub> between 2500 and 5000 uatm are relevant.

The most consistent impacts across larval yellowfin early life history processes occurred at the highest pCO<sub>2</sub> levels tested (>8500 uatm; Table 1). These levels are outside those predicted to occur in the next 300 years. However, there was evidence of significantly reduced survival at mean pCO<sub>2</sub> levels of  $\geq 4700$  uatm (experiment 1) and significantly reduced larval size (experiment 1) and extended egg hatch time (experiment 2) at mean pCO<sub>2</sub> levels  $\geq 2200$  uatm, that are relevant to near future predicted levels (Table 1). Significant effects at near future pCO<sub>2</sub> levels were not, however, consistently predicted in both trials.

We observed reduced survival of yellowfin larvae after 7 days of feeding in the first trial with increasing mean pCO<sub>2</sub>. The mean survival at the control level (pH = 8.23, pCO<sub>2</sub> = 368) was not significantly different to survival at pH 7.56 (pCO<sub>2</sub> ~2108) but was significantly greater than survival at pH 7.35 (pCO<sub>2</sub> ~4732) and pH 6.90 (pCO<sub>2</sub> ~8847). The relationship between survival and mean pCO<sub>2</sub> in the second trial was neither clear nor statistically significant, due in large part to the high level of intra-treatment variability in survival. There was a sudden mortality event (crash) in tank 15 (pH 8.1) on the last night of trial 2, associated with a very high density of larvae throughout the trial in the same tank. Such events are uncommon but have been observed previously in

Table 1. Experimental results for the pilot study on how ocean acidification may impact yellowfin tuna. Statistically significant differences between control (current day pCO<sub>2</sub> levels) and treatment levels are presented as red squares. The green squares represent where statistically significant effects were not detected in experiments 1 and 2. The mean pH and pCO<sub>2</sub> for the control tanks for experiments 1 and 2 were 8.2/368 and 8.1/464 respectively.

	Experiment 1			Experiment 2		
	pH range pCO <sub>2</sub> range			pH range pCO <sub>2</sub> range		
	7.6 2108	7.4 3820	6.9 9824	7.6 1719	7.2 4733	7.0 8847
Egg stage duration						
Survival						
Growth dry weight						
Growth standard length						

high larval density tanks at Achotines after 4–5 days of feeding. High survival in that tank, if it had been included in the analysis, would have provided a declining survival trend with increasing pCO<sub>2</sub> in trial 2.

The egg stage and growth results were more consistent between the two trials. Pair-wise comparisons for both experiments indicated a significant difference in dry weights between larvae reared in the control treatment (mean pCO<sub>2</sub> ~368 in trial 1 and ~464 in trial 2) and the highest pCO<sub>2</sub> treatments (mean pCO<sub>2</sub> ~9824 in trial 1 and ~8847 in trial 2) but not between larvae reared at the control and intermediate pCO<sub>2</sub> levels. Effects at intermediate pCO<sub>2</sub> levels were detected for standard length. Pair-wise comparisons for experiment 1 indicated that larvae reared in the treatment tanks with mean pCO<sub>2</sub> ~2108 and mean pCO<sub>2</sub> ~4732 were significantly smaller than those reared in the control treatment (pCO<sub>2</sub> ~368). Slower larval growth during the first week of feeding may subsequently impact survival by affecting foraging success and by prolonging stage durations, increasing larval size-dependent susceptibility to predation (review in Leggett and DeBlois 1994).

A significant positive relationship between mean pCO<sub>2</sub> and hours until complete hatching was detected in both experiments. The hatch times at mean pCO<sub>2</sub> values less than ~2200 were very similar in experiment 1, and the significant increasing trend in hatch time for this experiment represents an increase in mean hatch time at mean pCO<sub>2</sub> > ~8800 (target pH 6.9) as compared to the mean hatch times at the lower pCO<sub>2</sub> levels. In experiment 2 the mean hatch time at pH 8.1 was significantly less than that at each of pH 7.3, 7.7, and 6.9. The mean hatch time

at pH 7.3 was also significantly less than that at pH 6.9 in this experiment. The difference in hatch time between the current conditions and mean pCO<sub>2</sub> > ~8800 was one hour. Similar delays in hatching have also been observed during adverse physical conditions of very low dissolved oxygen levels and extremely high water temperatures (Wexler et al. 2011).

The purpose of the pilot study was to trial techniques to help design more intensive experimental trials. Power analyses of variability in survival responses within and between tanks/treatments indicated that more replicate tanks per treatment are needed to increase the statistical power and sensitivity of the experiments, so that the functional form of survival relationships can be identified and described. This was less of an issue for the growth analyses for which sample numbers were an order of magnitude higher. An additional design consideration for future experiments is the application of elevated pCO<sub>2</sub> conditions to the parental stock. Recent research has demonstrated that exposure of broodstock to elevated pCO<sub>2</sub> prior to spawning may reduce or remove negative impacts of elevated pCO<sub>2</sub> upon behavior of larvae derived from subsequent spawnings (Miller et al. 2012).

Future experimental trials should include a lower pCO<sub>2</sub> treatment (~1000 uatm) to test yellowfin larval responses to lower “near future” pCO<sub>2</sub> levels. Including the interactive effects of temperature and oxygen, which will also vary under future climate change (Gruber 2011) would also be beneficial. OA and other parameters such as temperature have already been shown to interact for some species (Nowicki et al. 2012; Munday et al. 2009a; Enzor

et al. 2013). Hypoxic zones in the ocean may increase in the future as a result of climate driven changes in temperatures and deep ocean mixing (Hofman and Schellhuber 2009). Survival during the onset of feeding in yellowfin tuna larvae is greatly affected by short-term oxygen deficits at water temperatures  $\geq 26$  °C (Wexler et al. 2011). Larval sensitivity to dO<sub>2</sub> might increase under elevated pCO<sub>2</sub> conditions, and this needs to be tested in future experiments.

The very high fecundity and relatively short generation time of yellowfin tuna may permit them to adapt more rapidly than less fecund, longer-lived species. It will be important to determine the extent to which inter-individual variation mediates different selection responses (Schlegel et al. 2012). Genetic analyses of samples taken from the current trials are in progress to assess whether some genotypes are more robust to changes in pCO<sub>2</sub>, and these results should provide insights on the designs needed to assess this important question. Results of our supplemental sampling on the feeding patterns, otolith morphology and histological condition of internal tissues of larvae will also permit us to further investigate how acidification affects early life stages. Analyses of samples taken from yolk-sac and early feeding larvae will allow us to assess at what point in development any effects on growth first became apparent.

The ultimate goal of this research is to provide information that will allow models such as the Spatial Ecosystem and Population Dynamics Model (SEAPODYM) to be parameterised to include acidification effects and subsequently enable scientists and tuna fishery managers to better understand how these changes in ocean chemistry will alter the distribution and abundance of yellowfin tuna. Environmental data are used in SEAPODYM to functionally characterise the habitat of the population depending on its thermal, biogeochemical, and forage preferences (Lehodey et al. 2008, 2010). To this end, it will be critical that further empirical trials are conducted to more clearly identify the functional form of the relationship between pH (or pCO<sub>2</sub>) and larval survival, and in particular, identify any interactions between pH (or pCO<sub>2</sub>) and other key physical oceanographic factors such as temperature and oxygen, so as to provide relevant information for appropriately altering the spawning-habitat index in SEAPODYM. Subsequent population level predictions of ocean acidification impacts will enhance the capacity of regional fisheries management organisations to make better-informed decisions regarding the management of the highly valuable tropical tuna resources, particularly with regard to attaining key sustainability-related objectives (Bell et al. 2013).

## References

- Allen B., Domenici P., McCormick M.I., Watson S. and Munday P. 2013. Elevated CO<sub>2</sub> affects predator-prey interactions through altered performance. *PLoS ONE* 8: e58520. doi:10.1371/journal.pone.0058520
- Barnett T.P., Pierce D.W., Achuta Rao K.M., Gleckler P.J., Santer B.D., Gregory J.M. and Washington W.M. 2005. Penetration of human-induced warming into the world's oceans. *Science* 309: 284–7.
- Baumann, H., Talmage S.C., and Gobler C.J. 2011. Reduced early life growth and survival in a fish in direct response to increased carbon dioxide. *Nature Climate Change*. doi:10.1038/NCLIMATE1291
- Bell J.D., Ganachaud A., Gehrke P.C., Griffiths S.P., Hobday A.J., Hoegh-Guldberg O., Johnson J.E., Le Borgne R., Lehodey P., Lough J.M., Matear R.J., Pickering T.D., Pratchett M.S., Sen Gupta A., Senina I. and Waycott M. 2013. Tropical Pacific fisheries and aquaculture will respond differently to climate change. *Nature Climate Change* 3: 591–599.
- Bignami S., Sponaugle S. and Cowen R.K. 2013a. Response to ocean acidification in larvae of a large tropical marine fish, *Rachycentron canadum*. *Global Change Biology* 19: 996–1006.
- Bignami S., Enochs I.C., Manzello D.P., Sponaugle S. and Cowen R.C. 2013b. Ocean acidification alters the otoliths of a pantropical fish species with implications for sensory function. *Proceedings of the National Academy of Sciences*. DOI 10.1073/pnas.1301365110.
- Bopp L., Resplandy L., Orr J.C., Doney S.C., Dunne J.P., Gehlen M., Halloran P., Heinze C., Ilyina T., Seferian R., Tjiputra J. and Vichi M. 2013. Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences* 10: 6225–6245. www.biogeosciences.net/10/6225/2013/doi:10.5194/bg-10-6225-2013
- Brauner C.J. and Baker D.W. 2009. Patterns of acid-base regulation during exposure to hypercarbia in fishes. p. 43–63. In: M.L. Glass and S.C. Woods (eds). *Cardio-respiratory control in vertebrates*.
- Briffa M., de la Haye K. and Munday P.L. 2012. High CO<sub>2</sub> and marine animal behaviour: Potential mechanisms and ecological consequences. *Marine Pollution Bulletin* 64: 1519–1528.
- Caldeira K. and Wickett M.E. 2003. Anthropogenic carbon and ocean pH. *Nature* 425: 365.

- Caldeira K. and Wickett M.E. 2005. Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *Journal of Geophysical Research*. 110:n.p.
- Checkley D.M., Dickson A.G., Takahashi M., Radich J.A., Eisenkolf N. and Asch R. 2009. Elevated CO<sub>2</sub> enhances otolith growth in young fish. *Science* 324:1683.
- Cripps I.C., Munday P.L. and McCormick M.I. 2011. Ocean acidification affects prey detection by a predatory reef fish. *PLoS ONE* 7: e22736. doi:10.1371/journal.pone.0022736
- Devine B.M., Munday P.L. and Jones G.P. 2012. Rising CO<sub>2</sub> concentrations affect settlement behaviour of larval damselfishes. *Coral Reefs* 31(1): 229–38.
- Devine B.M., Munday P.L. and Jones G.P. 2010. Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues. *Ecology Letters* 13:68–75.
- Domenici P., Allan B., McCormick M.I. and Munday P.L. 2012. Elevated CO<sub>2</sub> affects behavioral lateralization in a coral reef fish. *Biology Letters* doi:10.1098/rsbl.2011.0591.
- Enzor L., Zippay M.L. and Place S.P. 2013. High latitude fish in a high CO<sub>2</sub> world: Synergistic effects of elevated temperature and carbon dioxide on the metabolic rates of Antarctic notothenioids. *Comparative Biochemistry and Physiology, Part A* 164: 154–161.
- Esbaugh A.J., Heuer R. and Grosell M. 2012. Impacts of ocean acidification on respiratory gas exchange and acid-base balance in a marine teleost, *Opsanus beta*. *Journal of Comparative Physiology Part B* 182: 921–934
- Fabry V.J., Seibel B.A., Feely R.A., and Orr J.C. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes—ICES. *Journal of Marine Sciences* 65: 414–32.
- Feely R.A., Sabine C.L., Lee K., Berelson W., Kleypas J., Fabry V.J. and Millero F.J. 2004. Impact of anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> System in the oceans. *Science* 305: 362–6.
- Ferrari M.C.O., McCormick M.I., Munday P.L., Meekan M.G., Dixon D.L., Lonnstedt O. and Chivers D.P. 2012a. Effects of ocean acidification on visual risk assessment in coral reef fishes. *Functional Ecology* 26: 553–558; doi: 10.1111/j.1365-2435.2011.01951.x
- Ferrari M.C.O., Manassa R., Dixon D.L., Munday P.L., McCormick M.I., Meekan M.G., Sih A. and Chivers D.P. 2012b. Effects of ocean acidification on learning in coral reef fishes. *PLoS ONE* 7: e31478. doi:10.1371/journal.pone.0031478
- Franke A. and Clemmesen C. 2011. Effect of ocean acidification on early life stages of Atlantic herring (*Clupea harengus* L.). *Biogeosciences Discuss* 8: 7097–7126.
- Frommel A.Y., Stiebens V., Clemmesen C. and Havenhand J. 2010. Effect of ocean acidification on marine fish sperm (Baltic cod: *Gadus morhua*). *Biogeosciences*, 7: 3915–3919.
- Frommel A.Y., Maneja R., Lowe D., Malzahn A.M., Gefen A.J., Folkvord A., Piatkowski U., Reusch T.B.H. and Clemmesen C. 2011. Severe tissue damage in Atlantic cod larvae under increasing ocean acidification. *Nature Climate Change*, 2: 42–46. DOI: 10.1038/NCLIMATE1324
- Frommel A.Y., Schubert A., Piatkowski U. and Clemmesen C. 2012. Egg and early larval stages of Baltic cod, *Gadus morhua*, are robust to high levels of ocean acidification. *Marine Biology* DOI 10.1007/s00227-011-1876-3
- Gillett R. 2009. Fisheries in the economies of Pacific Island countries and territories. *Pacific Studies Series*, Asian Development Bank, Manila, Philippines.
- Gruber N. 2011. Warming up, turning sour, losing breath: ocean biogeochemistry under global change. *Philosophical Transactions of the Royal Society a-Mathematical Physical and Engineering Sciences* 369: 1980-1996 doi 10.1098/rsta.2011.0003
- Hofman M. and Shellhuber H. 2009. Oceanic acidification affects marine carbon pump and triggers extended marine oxygen holes. *Proceedings of the National Academy of Sciences* 106: 3017-3022; www.pnas.org/cgi/doi/10.1073/pnas.0813384106
- Houde E. 1989. Subtleties and episodes in the early life of fishes. *Journal of Fish Biology* 35: 29–38.
- Hurst T.P., Fernandez E.R., Mathis J.T., Miller J.A., Stinson C.M. and Ahgeak E. F. 2012. Resiliency of juvenile walleye pollock to projected levels of ocean acidification. *Aquatic Biology* 17: 247–259.
- Hurst T.P., Fernandez E.R. and Mathis J.T. 2013. Effects of ocean acidification on hatch size and larval growth of walleye pollock (*Theragra chalcogramma*). *ICES Journal of Marine Science* 70: 812-822. doi: 10.1093/icesjms/fst053

- Ilyina T., Zeebe R.E., Maier-Reimer E. and Heinze C. 2009. Early detection of ocean acidification effects on marine calcification. *Global Biogeochemical Cycles* 23: GB1008. doi:10.1029/2008GB003278.
- Ilyina T., Six K.D., Segsneider J., Maier-Reimer E., Li H. and Núñez-Riboni I. 2013. The global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth System Model in different CMIP5 experimental realizations. *Journal of Advances in Modeling Earth Systems*, doi: 10.1002/jame.20017, 2013.
- IPCC. 2007. *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II, and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* [Core Writing Team: Pachauri, R.K. and A. Reisinger, eds.] Geneva, Switzerland: IPCC.
- IPCC. 2011. *Workshop Report of the Intergovernmental Panel on Climate Change Workshop on Impacts of Ocean Acidification on Marine Biology and Ecosystems* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, K.J. Mach, G.-K. Plattner, M.D. Mastrandrea, M. Tignor and K.L. Ebi (eds.)]. IPCC Working Group II Technical Support Unit, Carnegie Institution, Stanford, California, United States of America, pp. 164.
- Jonz M.G. and Nurse C.A. 2006. Ontogenesis of oxygen chemoreception in aquatic vertebrates. *Respiratory Physiology and Neurobiology* 154: 139–152.
- Leggett W.C. and DeBlois E. (1994). Recruitment in marine fishes - is it regulated by starvation and predation in the egg and larval stages. *Netherlands Journal of Sea Research* 32: 119–134.
- Kikkawa T., Ishimatsu A. and Kita J. 2003: Acute CO<sub>2</sub> tolerance during the early developmental stages of four marine teleosts. *Environmental Toxicology* 18: 375–382.
- Lehodey P., Senina I. and Murtugudde R. 2008. A spatial ecosystem and populations dynamics model (SEAPODYM) modelling of tuna and tuna-like populations. *Progress in Oceanography* 78: 304–18.
- Lehodey P., Senina I., Sibert J., Bopp L., Calmettes B., Hampton J. and Murtugudde R. 2010. Preliminary forecasts of population trends for Pacific big-eye tuna under the A2 IPCC Scenario. *Progress in Oceanography* doi:10.1016/j.pocean.2010.04.021.
- Lehodey P., Senina I., Calmettes B., Hampton J., Nicol S. 2013. Modelling the impact of climate change on Pacific skipjack tuna population and fisheries. *Climatic Change* 119: 95-109. DOI 10.1007/s10584-012-0595-1,
- Margulies D., Suter J.M., Hunt S.L., Olson R.J., Scholey V.P., Wexler J.B. and Nakazawa A. 2007. Spawning and early development of captive yellowfin tuna (*Thunnus albacares*). *Fisheries Bulletin* 105: 249–65.
- Miller G.M., Watson S.A., Donelson J.M., McCormick M.I. and Munday P.L. 2012a. Parental environment mediates impacts of increased carbon dioxide on a coral reef fish. *Nature Climate Change*. DOI: 10.1038/NCLIMATE1599
- Munday P.L., Donelson J.M., Dixon D.L., and Endo G.G.K. 2009a. Effects of ocean acidification on the early life history of a tropical marine fish. *Proceedings of the Royal Society London B* 276: 3275–83.
- Munday P.L., Dixon D.L., Donelson J.M., Jones G.P., Pratchett M.S., Devitsina G.V. and Døving K.B. 2009b. Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proceeding of the National Academy of Sciences* 106: 1848–1852.
- Munday P.L., Crawley N. and Nilsson G.E. 2009c. Interacting effects of elevated temperature and ocean acidification on the aerobic performance of coral reef fishes. *Marine Ecology Progress Series* 388: 235–242.
- Munday P.L., Gagliano M., Donelson J.M., Dixon D.L. and Thorrold S. 2011a. Ocean acidification does not affect the early life history development of a tropical marine fish. *Marine Ecology Progress Series* 423: 211–221. doi: 10.3354/meps08990
- Munday P.L., Hernaman V., Dixon D.L. and Thorrold S.R. 2011b. Effect of ocean acidification on otolith development in larvae of a tropical marine fish. *Biogeosciences* 8: 1631–1641; doi:10.5194/bg-8-1631-2011
- Nilsson G., Dixon D.L., Domenici P., McCormick M.I., Sorensen C., Watson S. and Munday P.L. 2012. Near future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function. *Nature Climate Change*. doi:10.1038/NCLIMATE1352.
- Nowicki J.P., Miller G.M. and Munday P.L. 2012. Interactive effects of elevated temperature and CO<sub>2</sub> on foraging behavior of juvenile coral reef fish. *Journal of Experimental Marine Biology and Ecology* 412: 46–51. doi.org/10.1016/j.jembe.2011.10.020
- Orr J.C., Fabry V.J., Aumont O., Bopp L., Doney S.C., Feely R.A., Gnanadesikan A., et al. 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437: 681–6.

- Pelster B. 2008. Gas exchange. p. 102–103. In: Finn R.N. and Kapoor B.G. (Eds.), *Fish Larval Physiology, Part 2, Respiration & Homeostasis*. Science Publishers, Enfield, New Hampshire, United States of America.
- Raven J., Caldeira K., Elderfield H., Hoegh-Guldberg O., Liss P., Riebesell U., Shepherd J., Turley C. and Watson A. 2005. Ocean acidification due to increasing atmospheric carbon dioxide. Policy Document 12/05. London: Royal Society London.
- Riebesell U., Fabry V.J., Hansson L. and Gattuso J.-P. (eds.). 2010. *Guide to best practices for ocean acidification research and data reporting*. Luxembourg: Publications Office of the European Union.
- Sabine C.L., Feely R.A., Gruber N., Key R.M., Lee K., Bullister J.L. et al. 2004. The oceanic sink for CO<sub>2</sub>. *Science* 305:367–71.
- Schaefer K.M. 2001. Reproductive biology of tunas. p. 225–270. In: B. A. Block and E. D. Stevens (eds). *Fish physiology*, Vol. 19, Tuna: physiology, ecology, and evolution. Academic Press, San Diego, California, United States of America.
- Schlegel P., Havenhand J.N., Gillings M.R. and Williamson J.E. 2012. Individual variability in reproductive success determines winners and losers under ocean acidification: A case study with sea urchins. *PLoS ONE* 7:e53118
- Simpson S.D., Munday P.L., Wittenrich M.L., Manassa R., Dixon, D.L., Gagliano M. and Yan, H.Y. 2011. Ocean acidification erodes crucial auditory behaviour in a marine fish, *Biology Letters* 7: 917–920, DOI:10.1098/rsbl.2011.0293
- Sundin J., Rosenqvist G. and Berglund A. 2013. Altered oceanic pH impairs mating propensity in a pipefish. *Ethology* 119: 86–93.
- Tseng Y.-C., Yu M.H., Stumpp M., Yin L.-Y., Melzner F. and Hwang P.-P. 2013. CO<sub>2</sub>-driven seawater acidification differentially affects development and molecular plasticity along life history of fish (*Oryzias latipes*). *Comparative Biochemistry and Physiology Part A* 165: 119–130.
- Wexler J.B., Margulies D., and Scholey V.P. 2011. Temperature and dissolved oxygen requirements for survival of yellowfin tuna, *Thunnus albacares*, larvae. *Journal of Experimental Marine Biology and Ecology* 404: 63–72.
- Williams P. and Terawasi P. 2009. Overview of tuna fisheries in the Western and Central Pacific Ocean, including economic conditions—2008. WCPFC-SC5-2009-GN-WP-01. Western and Central Pacific Fisheries Commission Scientific Committee Fifth Regular Session, 10–21 August 2009, Port Vila, Vanuatu.

## Author affiliations

1. Vernon Scholey is the Director of Achotines Laboratory (Inter-American Tropical Tuna Commission), Las Tablas, Los Santos, Panama.
2. Don Bromhead, Simon Hoyle and Simon Nicol are Fisheries Scientists with the Oceanic Fisheries Programme, Secretariat of the Pacific Community, BP D5, 98848 Noumea CEDEX, New Caledonia.
3. Daniel Margulies, Jeanne Wexler, Maria Santiago and Cleridy Lennert-Cody are Fisheries Scientists with the Inter-American Tropical Tuna Commission, 8604 La Jolla Shores Drive, La Jolla, California, 92037, USA.
4. Jane E. Williamson is a Senior Lecturer within Biological Sciences and runs the Marine Ecology Group at Macquarie University, NSW, 2109, Australia.
5. Jon Havenhand is a larval ecologist working on ocean acidification in the Tjärnö Marine Biological Laboratory, University of Gothenburg, 45296 Strömstad, Sweden.
6. Tatiana Ilyina leads the Ocean Biogeochemistry Group at the Max Planck Institute for Meteorology, Bundesstrasse 53, 20146 Hamburg, Germany.
7. Patrick Lehodey is head of the Marine Ecosystem Department of the Space Oceanography Division of CLS, France, 8-10 rue Hermes, 31520, Ramonville St Agne, France.

© Copyright Secretariat of the Pacific Community, 2014

All rights for commercial / for profit reproduction or translation, in any form, reserved. SPC authorises the partial reproduction or translation of this newsletter for scientific, educational or research purposes, provided that SPC and the source document are properly acknowledged. Permission to reproduce the document and/or translate in whole, in any form, whether for commercial / for profit or non-profit purposes, must be requested in writing.

Original SPC artwork may not be altered or separately published without permission.

The views expressed in this Bulletin are those of the authors and are not necessarily shared by the Secretariat of the Pacific Community.

Original text: English

Secretariat of the Pacific Community, Fisheries Information Section, BP D5, 98848 Noumea Cedex, New Caledonia

Telephone: +687 262000; Fax: +687 263818; cfpinfo@spc.int; <http://www.spc.int/fame>